

| Product | Dosage / Volume / Unit of Measurement |
|---|--|
| First aid kits | |
| Paracetamol | 500 mg |
| Meloxicam | |
| Individual sterile dressing package with elastic first aid compression component with protective moisture-resistant shell | |
| Sterile gauze bandages | |
| Chemical remedy to stop bleeding | |
| Mechanical means for a stop of bleeding like "CAT" | |
| Gel-based occlusive thoracic bandage | |
| Non-woven adhesive plaster | |
| Nitrile non-sterile medical examination gloves | |
| Nasopharyngeal airway (airway, tube) with lubricant | |
| Atropine | injection: 1 mg (sulphate) in ampoules of 1 ml |

| | |
|--|--|
| Heparin sodium | injections: 5000 IU / ml |
| Hydroxyethylstarch | 6% infusion solution (60 mg / ml) |
| Hydroxyethylstarch | 10% solution for infusion (100 mg / ml) |
| Dexamethasone | injection: 4 mg / ml 1 ml in ampoules (as disodium phosphate salt) |
| Diazepam | injections: 5 mg / ml |
| Dopamine | injection: 40 mg / ml (hydrochloride) 5 ml in ampoules |
| Epinephrine /Adrenaline | injection: 1 mg (as hydrochloride and hydrotartrate) 1 ml in ampoules, corresponding to 1.82 mg of adrenaline tartrate in 1 ml |
| Ethanol | solution: 70% (denatured) |
| Isosorbide dinitrate | solution for infusion: 1 mg / ml |
| Insulin injection (fast-acting) | injection: 40 IU / ml in 10 ml vials |
| Insulin injection (fast-acting) | injections of 100 IU / ml in 5 ml; 10 ml in a bottle |
| Sodium lactate, complex solution of Sodium chloride + Potassium chloride + Calcium chloride + Sodium lactate | injection of 100 U / ml and 5 ml; 10 ml in a bottle |
| Mannitol | solution for infusion: 15% 100 ml, 200 ml, 400 ml in bottles, 100 ml, 250 ml, 500 ml in containers |

| | |
|---|--|
| Naloxone | injection: 400 mcg (hydrochloride) 1 ml in ampoules |
| Sodium hydrogen carbonate | injection: 1,4 % - 8,4 % |
| Sodium chloride | solution for injection / infusion: 0.9% isotonic (equivalent to Na + 154 mmol / l, Cl - 154 mmol / l) |
| Complex solution of Sodium chloride + Potassium chloride + Calcium chloride | solution for infusion |
| Neostigmine | injections: 0.5 mg / ml in 1 ml ampoules |
| Neostigmine | injections of 2.5 mg (methyl sulfate) in 1 ml ampoules |
| Ondansetron | injection: 2 mg / ml 2 ml in ampoules (as hydrochloride) |
| Ondansetron | injection of 2 mg / ml 4 ml in ampoules (as hydrochloride) |
| Povidone-iodine | solution: 10% (equivalent to 1% active iodine) |
| Salbutamol | metered dose inhaler (aerosol): 100 mcg (as sulfate) per 1 dose |
| Furosemide | injections: 10 mg / ml in 2 ml ampoules |
| Chlorhexidine | solution: 0.05% (bigluconate) |
| Tranexamic acid | injection: 50 mg / ml in an ampoule |

| | |
|---------------------|---|
| Tranexamic acid | 100 mg / ml in ampoules |
| Fresh-frozen plasma | doses |
| Red blood cells | doses |
| Cryoprecipitate | doses |
| Oxygen | gas (liquid) |
| Oxygen | gas |
| Sevoflurane | liquid for inhalation |
| Ketamine | injections: 50 mg / ml 2 ml injections: 50 mg / ml 10 ml injections: 10 mg / ml 10 ml injections: 10 mg / ml 50 ml |
| Propofol | injections: 10 mg / ml 100 ml injections: 20 mg / ml |
| (Thiopental) | injections: 0.5 g injections: 1 g |
| Bupivacaine | injections: 5 mg / ml 4 ml injections: 5 mg / ml 5 ml injections: 5 mg / ml 10 ml injections: 5 mg / ml 20 ml |
| Bupivacaine | 0.5% (hydrochloride) 4 ml in ampoules |

| | |
|-------------------------------|--|
| Lidocaine | injection: 1% 2 ml in ampoules |
| Lidocaine | 2% injection (hydrochloride) of 2 ml in ampoules |
| Lidocaine | 5% (hydrochloride) 2 ml in ampoules |
| Morphine | injection: 10 mg (sulphate or hydrochloride) 1 ml in ampoules |
| Fentanyl | solution for injection: 0.05 mg / ml in 2 ml ampoules |
| Acetylcysteine | injections: 100 mg / ml 3 ml in ampoules injections: 200 mg / ml in 10 ml ampoules |
| Amoxicillin + Clavulanic acid | tablets: 500 mg / 125 mg; powder for injection: 1000 mg + 200 mg |
| Ceftriaxone | powder for solution for injection: 1 g (as sodium salt) in a vial powder for solution for injection: 1000 mg (as sodium salt) in a vial |
| Imipenem + Cilastatin | powder for solution for injection: 500 mg (as monohydrate) + 500 mg (as sodium salt) in vials |
| Meropenem | powder for injection / infusion: 1 g (as trihydrate) |
| Daptomycin | powder for concentrate for solution for infusion: 500 mg powder for solution for infusion: 500 mg |
| Linezolid | solution for infusion: 2 mg / ml of 300 ml |
| Cefepime | powder for solution for injection: 1000 mg; |

| | |
|--------------------|--|
| Azithromycin | solid oral dosage form: 500 mg solution for oral use: 200 mg / 5 ml |
| Ciprofloxacin | solution for intravenous infusion: 2 mg / ml (as hyclate) tablets: 500 mg (as hydrochloride) |
| Metronidazole | injection: 500 mg per 100 ml vial |
| Vancomycin | powder for solution for injection: 1000 mg (hydrochloride) in a vial |
| Flucytosine | solution for infusion: 2.5 g / 250 ml |
| Protamine sulfate | injections: 10 mg / ml (10,000 IU) in 5 ml |
| Glucose) | solution for injection: 5% (isotonic) solution for injection: 40 % |
| Potassium chloride | 0,04 |
| Potassium chloride | 0,075 |
| Suxamethonium | 50 mg (chloride) / ml 2 ml in ampoules 20 mg / ml 5 ml in ampoules (suxamethonium iodide) powder for solution for injection (chloride) in a vial |
| Atracurium | 10 mg / ml (besylate) |
| Vecuronium | powder for solution for injection: 10 mg (bromide) |

| | |
|--|---|
| Enoxaparin | 20 mg / 0.2 ml; 40 mg / 0.4 ml; 100 mg / 1 ml; 150 mg / 1 ml |
| Metoprolol tartrate | 1 mg / ml 5 ml |
| Phenylephrine hydrochloride | 1% 1 ml |
| Norepinephrine Tartrate | 2 mg / ml 4 ml 2 mg / ml 8 ml |
| Stomach wash kit | |
| Set for catheterization of the bladder | |
| Traction tire | |
| Portable oxygen cylinders | |
| Central vein catheterization kits | |
| Set for intraosseous access | |
| Kits for lumbar puncture | |
| Tracheostomy set | |

| | |
|--|--|
| Tracheal intubation kit | |
| Set for drainage of the pleural cavity | |
| Immobilization tires of various types and sizes | |
| Syringes with needles (different volumes) | |
| Catheters for peripheral veins (different sizes) | |
| System for transfusion of infusion solutions | |
| Non - sterile bandage (different sizes) | |
| Elastic fixing bandage (different sizes) | |
| Sterile material for anti-burn measures | |
| Sterile material for wound surface treatment | |
| Hydrogel anti-burn bandage of different sizes | |
| External fixation device | |
| Gel-based occlusive thoracic bandage (sticker) (with or without valve) | |

| | |
|---|--|
| Scissors for cutting clothes and shoes (atraumatic) | |
| Film (valve) for artificial lung ventilation | |
| Puncture needle for decompression of the pleural cavity | |
| Hard neck collar with size adjustment | |
| Waterproof marker | |
| Intraosseous catheters (FAST, B.I.G, EASY IO) | |
| Eyeshield | |
| Reinforced tape | |
| Water gel burn dressings | |
| Trauma shears or sling cutter | |
| Occlusive film | |
| Hemostatic dressings. Verified producers: Krovospas (local), Combat Gauze quick dot, Celox gauze, Chito Gauze | |
| Hemostatic dressing of quick dot combat type | |

| |
|--|
| Institute of Pediatrics, Obstetrics and Gynecology of the National Academy of Medical Sciences of Ukraine |
| Disinfection tablets/pills for drinking water |
| Skin barrier film 3M Cavilon |
| Disposable skin clips |
| Tachcomb 9,5/4,8 & 4,8/4,8 |
| Multipurpose manual resuscitation bag, for adults |
| Hemostatic Sponges |
| Gel bandages for burn and wound treatment |